

## A new role for the extracellular calcium-sensing receptor demonstrated by using CCK-eGFP BAC mice

R. John MacLeod

Department of Physiology/Gastrointestinal Disease Research Unit, Queen's University, Kingston, Ontario, Canada

Submitted 28 January 2011; accepted in final form 9 February 2011

ONE OF THE THINGS THAT MAKE studying gastrointestinal processes both exciting and fun is the application of new techniques to address a problem where understanding has been constrained by previous techniques. Two complementary papers (10, 19) in this issue illustrate such an application. The new technique is the use of bacterial artificial chromosome (BAC) transgenic mice. The problem is nutrient sensing by enteroendocrine cells. Widely used in neuroscience research (5), the BAC mice allow either targeted gene deletion or overexpression (20). Many transgenic BAC mice express regulatory sequences together with enhanced green fluorescent protein (eGFP), which also allows identification of specific cell types. The eGFP allows these cells to be recovered using FACS for further analysis. Such BAC transgenic mice may then be mated with knockout or reporter mice to further define the presence of either a receptor or a pathway activated by the receptor.

The satiating effects of dietary protein may be due in part to cholecystokinin (CCK) secretion from duodenal I cells. Although aromatic amino acids have been understood to be mediators of CCK secretion from these cells, the technical difficulties of recovering sufficient amounts of homogeneous I cells from fetal or adult intestine are character building. The use of cell lines (i.e., STC-1) that are tumorigenic but endocrine derived is informative but equivocal. Consequently, little is understood about the cellular mechanism by which amino acids affect CCK secretion. The extracellular calcium-sensing receptor (CaSR), originally cloned from the parathyroid, but found in distinct regions of the gastrointestinal system, may be activated by aromatic amino acids, consistent with the CaSR being an amino acid sensor (1, 2). Because the CaSR knockout is perinatal lethal, it was “rescued” by taking parathyroid hormone (PTH) out of the system through breeding with CaSR<sup>+/-</sup>PTH<sup>+/-</sup> heterozygotes. The resulting CaSR<sup>-/-</sup>PTH<sup>-/-</sup> are sustained with 2% Ca<sup>2+</sup> in their drinking water (8, 9). Analyses using heterologous CaSR expressing cells have shown differences in intracellular Ca<sup>2+</sup> oscillation frequency (13, 14) generated by aromatic amino acids compared with other ligands, but direct *in vivo* evidence from enteroendocrine cells has been lacking. Indeed, we still don't understand the physiological importance of circulating amino acids in regulating the CaSR's activity outside the gastrointestinal tract (3).

The study by Liou et al. (10) from the laboratories of Raybould and Wank is as impressive as it is elegant. They used FACS to isolate and enrich duodenal I cells from the CCK-eGFP BAC transgenic mice and demonstrated that the CaSR

was present on these cells. Expression of other putative mediators of amino acid sensing, such as PepT1 or the umami taste receptor, were also assessed. Liou et al. show that these cells generated intracellular Ca<sup>2+</sup> ([Ca<sup>2+</sup>]<sub>i</sub>) changes that were CaSR dependent, stereoselective for L-phenylalanine (L-Phe), and responsive to type II calcimimetic cincalcet. Increases in extracellular Ca<sup>2+</sup> ([Ca<sup>2+</sup>]<sub>o</sub>) did not stimulate CCK secretion from eGFP I cells, but L-Phe stimulation of CCK required “physiological” levels of calcium, consistent with how L-amino acids allosterically activate the CaSR. Liou et al. also ablated the CaSR by mating the CCK-eGFP mice with CaSR/PTH double homozygous knockout mice. Interestingly, they found that [Ca<sup>2+</sup>]<sub>o</sub> stimulations and both L-Phe and cincalcet caused a decrease in CCK secretion compared with basal secretion. Particularly cogent are their comments reviewing CaSR-mediated G<sub>q</sub> coupling with L-Phe stimulating G<sub>12/13</sub> and the cation channel TRPC1. They suggest that the splice variant of the CaSR, which lacks exon 5 and is known to be present in the CaSR/PTH double homozygous knockout mice (12), may explain the effect of cincalcet reducing CCK secretion below basal levels, but clearly more work needs to be done. The same authors recently used CCK-eGFP BAC mice to demonstrate a role for GPR40 to mediate long-chain fatty acid-induced secretion of CCK (11).

Equally impressive and elegant is the study by Wang et al. (19) from the laboratory of Liddle. Wang et al. also used the CCK-eGFP BAC mice and isolated mucosal cells with FACS and demonstrated the CaSR's expression on them. They also measured changes in [Ca<sup>2+</sup>]<sub>i</sub> and showed that both L-Phe and tryptophan caused transient increases that were prevented by the CaSR inhibitor Calhex 231. Both amino acids stimulated CCK release, which was prevented by this inhibitor. Whole cell patch-clamp technique demonstrated the CCK-eGFP cells had a predominantly outwardly rectifying K<sup>+</sup> current; this was inhibited by addition of L-Phe resulting in the depolarization required for CCK secretion from these cells. Both this paper (19) and the study from Wank and colleagues (10) extend our understanding of CaSR in the gastrointestinal system and illustrate the complexity of nutrient sensing in enteroendocrine cells.

We will see greater use of eGFP BAC [usually generated from the GENSAT Brain Atlas Project (5) and obtained from Mutant Mouse Regional Resource Centers] mice to address problems that have heretofore been technically challenging. For example, a SOX9eGFP mouse, also generated from GENSAT, will allow further definition of intestinal epithelial stem cell niche determinants (6), and a gastrin eGFP BAC mouse has recently been characterized (18). It is important to note that BAC mice generated by the GENSAT project are on a CD1 background and consequently must be backcrossed on the same background as the investigators' transgenic or knockout mice to

Address for reprint requests and other correspondence: R. J. MacLeod, Canada Research Chair in Gastrointestinal Cell Physiology, Dept. of Physiology, Queen's Univ., Rm 3-003, GIDRU Wing, 76 Stuart St., Kingston, ON, Canada K7L 2V7 (e-mail: rjm5@queensu.ca).

permit quantitative comparisons. In general, demonstrating that FACS isolation itself has not influenced the function of the isolated cells is required. Some understanding about whether BAC reporters faithfully recapitulate expression of the endogenous gene may also be worthwhile. Although the dramatic and exciting studies using the NF- $\kappa$ B reporter mice deserve citation (4, 7), eGFP tags on specific gastrointestinal cell types such as the stem cell marker Lrg5 (Jackson Laboratories) allow experimental designs that previously were only dreamed about (16, 17). For example, gene microarray analysis of Lrg5-eGFP cells showed Ror2 (the coreceptor for Wnt5a) but not CaSR, yet CaSR expression is stimulated by Wnt5a (13; unpublished observations). Understanding Wnt coreceptor plasticity, signaling, and Wnt family molecule plasticity in the normal and diseased intestine, as well as nutrient sensing in enteroendocrine cells, will be facilitated by eGFP BAC reporter mice.

#### GRANTS

Our work was supported by an National Sciences and Engineering Research Council-Dairy Farmers of Canada Cooperative Grant, the Crohns and Colitis Foundation of Canada, and the Canada Research Chair Program.

#### DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the author(s).

#### REFERENCES

1. Brown EM, MacLeod RJ. Extracellular calcium sensing and extracellular calcium signaling. *Physiol Rev* 81: 239–297, 2001.
2. Conigrave AD, Quinn SJ, Brown EM. L-amino acid sensing by the extracellular Ca<sup>2+</sup>-sensing receptor. *Proc Natl Acad Sci USA* 97: 4814–4819, 2000.
3. Conigrave AD, Mun HC, Brennan SC. Physiological significance of L-amino acid sensing by extracellular Ca<sup>2+</sup>-sensing receptors. *Biochem Soc Trans* 35: 1195–1198, 2007.
4. Ding S, Chi MM, Scull BP, Rigby R, Schwerbrock NM, Magness S, Jobin C, Lund PK. High-fat diet: bacteria interactions promote intestinal inflammation which precedes and correlates with obesity and insulin resistance in mouse. *PLoS One* 5: e12191, 2010.
5. Gong S, Zheng C, Doughty ML, Losos K, Didkovsky N, Schambra UB, Nowak NJ, Joyner A, Leblanc G, Hatten ME, Heintz N. A gene expression atlas of the central nervous system based on bacterial artificial chromosomes. *Nature* 425: 917–925, 2003.
6. Gracz AD, Ramalingam S, Magness ST. Sox9 expression marks a subset of CD24-expressing small intestine epithelial stem cells that form organoids in vitro. *Am J Physiol Gastrointest Liver Physiol* 298: G590–G600, 2010.
7. Joo YE, Karrasch T, Muhlbauer M, Allard B, Narula A, Herfarth HH, Jobin C. Tomato lycopene extract prevents lipopolysaccharide-induced NF-kappaB signaling but worsens dextran sulfate sodium-induced colitis in NF-kappaBEGFP mice. *PLoS One* 4: e4562, 2009.
8. Kantham L, Quinn SJ, Egbuna OI, Baxi K, Butters R, Pang JL, Pollack MR, Goltzman D, Brown EM. The calcium-sensing receptor (CaSR) defends against hypercalcemia independently of its regulation of parathyroid hormone secretion. *Am J Physiol Endocrinol Metab* 297: E915–E923, 2009.
9. Kos CH, Karaplis AC, Peng JB, Hediger MA, Goltzman D, Mohammad KS, Guise TA, Pollack MR. The calcium-sensing receptor is required for normal calcium homeostasis independent of parathyroid hormone. *J Clin Invest* 111: 1021–1028, 2003.
10. Liou AP, Sei Y, Zhao X, Feng J, Lu X, Thomas C, Pechhold S, Raybould HE, Wank SA. The extracellular calcium sensing receptor is required for cholecystokinin secretion in response to L-phenylalanine in acutely isolated intestinal I cells. *Am J Physiol Gastrointest Liver Physiol*. doi:10.1152/ajpgi.00342.2010.
11. Liou AP, Lu X, Sei Y, Zhao X, Perchold S, Carrero RJ, Raybould HE, Wank SA. The G-protein-coupled receptor GPR40 directly mediates long-chain fatty acid-induced secretion of cholecystokinin. *Gastroenterology* Oct 16 2010. [Epub ahead of print]
12. Oda Y, Tu CL, Chang W, Crumrine D, Komuves L, Mauro T, Elias PM, Bikle DD. The calcium-sensing receptor and its alternatively spliced form in murine epidermal differentiation. *J Biol Chem* 275: 1183–1190, 2000.
13. Pacheco II, MacLeod RJ. CaSR stimulates secretion of Wnt5a from colonic myofibroblasts to stimulate CDX2 and sucrase-isomaltase using Ror2 on intestinal epithelia. *Am J Physiol Gastrointest Liver Physiol* 295: G748–G759, 2008.
14. Rey O, Young SH, Papazyan R, Shapiro MS, Rozengurt E. Requirement of the TRPC1 cation channel in the generation of transient Ca<sup>2+</sup> oscillations by the calcium-sensing receptor. *J Biol Chem* 281: 38730–38737, 2006.
15. Rey O, Young SH, Yuan J, Slice L, Rozengurt E. Amino acid-stimulated Ca<sup>2+</sup> oscillations produced by the Ca<sup>2+</sup>-sensing receptor are mediated by a phospholipase C/inositol 1,4,5-trisphosphate independent pathway that requires G12, Rho, filamin-A and the actin cytoskeleton. *J Biol Chem* 280: 22875–22882, 2005.
16. Sato T, Vries RG, Snippert HJ, van der Wetering M, Barker N, Stange DE, van Es J, Abo A, Kujala P, Peters P, Clevers H. Single Lrg5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature* 459: 262–265, 2009.
17. Snippert HJ, Clevers H. Tracking adult stem cells. *EMBO Rep* 12: 113–122, 2011.
18. Takaishi S, Shibata W, Tomita H, Jin G, Yang X, Ericksen R, Dubaykovskaya Z, Asfaha S, Quante M, Betz KS, Shulkes A, Wang TC. In vivo analysis of mouse gastrin gene regulation in enhanced GFP-BAC transgenic mice. *Am J Physiol Gastrointest Liver Physiol* 300: G334–G344, 2011.
19. Wang Y, Chandra R, Samsa LA, Gooch B, Fee BE, Cook MJ, Vigna SR, Grant AO, Liddle RA. Amino acids stimulate cholecystokinin release through the calcium-sensing receptor. *Am J Physiol Gastrointest Liver Physiol*. doi:10.1152/ajpgi.00387.2010.
20. Yang XD, Ai W, Asfaha S, Bhagat G, Friedman RA, Jin G, Park H, Shykind B, Diacovo TG, Falus A, Wang TC. Histamine deficiency promotes inflammation-associated carcinogenesis through reduced myeloid maturation and accumulation of CD11b(+)LyG(+) immature myeloid cells. *Nat Med* 17: 87–95, 2011.